

Identification of *Tinomiscium petiolare* from Vietnam using the DNA barcode

DOI: 10.30901/2227-8834-2021-2-114-122

УДК 582:575:581.9

Поступление/Received: 16.01.2021

Принято/Accepted: 12.05.2021



В. В. ТХИНЬ^{1*}, Р. В. ДУДКИН^{1,2}, Л. Д. ЧАК³,
Н. В. ЧИНЬ³, Q. V. HOI⁴, N. T. LIEN³

Идентификация *Tinomiscium petiolare* из Вьетнама с помощью штрих-кода ДНК

Б. Б. ТХИНЬ^{1*}, Р. В. ДУДКИН^{1,2}, Л. Д. ЧАК³,
Х. В. ЧИНЬ³, К. В. ХОЙ⁴, Н. Т. ЛИЕН³

¹ School of Natural Sciences, Far Eastern Federal University,
10 Ajax Bay, Russky Island, Vladivostok 690922, Russia

² Botanical Garden Institute, Far-East Branch of the Russian
Academy of Sciences,
142 Makovskogo St., Vladivostok 690024, Russia

³ Faculty of Natural Sciences, Hong Duc University,
565 Quang Trung Street, Thanh Hoa 40130, Vietnam

⁴ Tay Nguyen Institute for Scientific Research,
Vietnam Academy of Science and Technology,
116 Xo Viet Nghe Tinh Street, Dalat 66107, Vietnam

*✉ buibaotinh9595@gmail.com

¹ Школа естественных наук,

Дальневосточный федеральный университет,
690922 Россия, г. Владивосток, о. Русский, п. Аякс, 10

² Ботанический сад-институт ДВО РАН,
690024 Россия, г. Владивосток, ул. Маковского, 142

³ Факультет естественных наук, Университет Хонгдык,
40130 Вьетнам, г. Тханьхоа, ул. Куанг Чунг, 565

⁴ Институт научных исследований Тай Нгуен
Вьетнамской академии наук и технологий,
66107 Вьетнам, г. Далам, ул. Со Вьет Нге Тинь, 116

*✉ buibaotinh9595@gmail.com

Background. *Tinomiscium petiolare* Hook.f. & Thomson is a medicinal species of the family Menispermaceae. This species is currently being intensively exploited for therapeutic purposes. Precise and rapid identification of *T. petiolare* is critical and essential for the classification, propagation, use and conservation of its genetic resources. In recent years, DNA barcoding has been known to be a fast and sensitive method for identifying species at any stage of development, using short DNA sequences. In this study we have performed the identification of *T. petiolare* specimens in Vietnam based on the sequence analysis of 4 DNA barcode loci: ITS, *matK*, *rbcl* and *rpoC*.

Materials and methods. Total DNA was extracted from leaf samples using DNeasy Plant Mini Kit. PCR amplification of the ITS, *matK*, *rbcl* and *rpoC* regions was carried out on the GeneAmp PCR System 9700 with specific primers. The purified PCR products were sequenced on the ABI 3500 Genetic Analyzer system, using BigDye® Terminator v3.1 Cycle Sequencing Kit. These genetic sequences were analyzed and compared, and a phylogenetic tree was constructed using BioEdit, BLAST, and MEGA 6 programs.

Results and conclusion. The success rate of amplification and sequencing was 100% for all 4 DNA barcode loci (ITS, *matK*, *rbcl* and *rpoC*) in the studied specimens. The produced sequence sizes of ITS, *matK*, *rbcl* and *rpoC* in the specimens were 574 bp, 810 bp, 527 bp and 488 bp, respectively. Further, we identified that all studied specimens were genetically related to each other and associated with the same species *T. petiolare*. Overall, the results of the study generated the most complete DNA barcode database of *T. petiolare* collected in Vietnam, contributing to the taxonomy and identification of this species.

Key words: Menispermaceae, sequence analysis, molecular identification, ITS, *matK*, *rbcl*, *rpoC*.

Актуальность. *Tinomiscium petiolare* Hook.f. & Thomson (тиномисциум черешковый) – лекарственный вид семейства Menispermaceae Juss. (Луносемянниковые). Этот вид в настоящее время активно используется в лечебных целях. Точная и быстрая идентификация *T. petiolare* имеет решающее значение для классификации, размножения, использования и сохранения его генетических ресурсов. В последние годы стало известно, что ДНК-штрихкодирование является быстрым и чувствительным методом идентификации видов на любой стадии развития с использованием коротких последовательностей ДНК. В этом исследовании мы провели идентификацию образцов *T. petiolare* во Вьетнаме на основе анализа последовательностей ДНК-штрихкодов 4 локусов: ITS, *matK*, *rbcl* и *rpoC*.

Материалы и методы. Тотальную ДНК экстрагировали из образцов листьев с помощью DNeasy Plant Mini Kit. ПЦР-амплификацию участков ITS, *matK*, *rbcl* и *rpoC* проводили в амплификаторе GeneAmp PCR System 9700 со специфическими праймерами. Очищенные продукты ПЦР секвенировали с помощью системы ABI 3500 Genetic Analyzer с использованием BigDye® Terminator v3.1 Cycle Sequencing Kit. Полученные последовательности были проанализированы, сравнены, и построено филогенетическое дерево с помощью программ BioEdit, BLAST и MEGA 6.

Результаты и выводы. Степень результативности амплификации и секвенирования составила 100% для ДНК-штрихкодов всех 4 локусов (ITS, *matK*, *rbcl* и *rpoC*) исследуемых образцов. Размеры последовательностей ITS, *matK*, *rbcl* и *rpoC* исследуемых образцов, которые мы получили, составляли 574 пн, 810 пн, 527 пн и 488 пн соответственно. Кроме того, мы определили, что все исследуемые образцы генетически связаны друг с другом и относятся к одному и тому же виду *T. petiolare*. В целом результаты исследования дали самую полную базу данных о ДНК-штрихкоде образцов *T. petiolare*, собранных во Вьетнаме, что способствовало идентификации и уточнению таксономии вида.

Ключевые слова: Menispermaceae, анализ последовательности, молекулярная идентификация, ITS, *matK*, *rbcl*, *rpoC*.

Introduction

Menispermaceae is a large family with about 68 genera and 440 species, including the genus *Tinomiscium* (Christenhusz, Byng, 2016). *Tinomiscium petiolare* is a medicinal species of the genus *Tinomiscium*. This species is found in China, India, Indonesia, Malaysia, Myanmar, Papua New Guinea, Philippines, Thailand, and Vietnam (Forman, 1988; Ho, 2000; Ghollasimood et al., 2012). The plants grow naturally and sparsely in mixed forests at elevations from 200 to 600 m. According to traditional medicine, *T. petiolare* has the effects on hemostasis, treatment of osteoarthritis pain, toothache, and cardiovascular disease (Van Valkenburg, Bunyapraphatsara, 2001; Chi, 2012).

Currently, *T. petiolare* is being intensively exploited for therapeutic purposes. Precise and rapid identification is critical for the classification, propagation, use and conservation of its genetic resources. Identification of this species is based on the morphological characteristics that have been studied (Ho, 2000; Chinh et al., 2015), but if the specimen is incomplete or crushed, it will be difficult to ensure high accuracy of morphological identification. Therefore, in order to overcome the disadvantages of morphology-based biological classification, the classification methods based on genetic materials have been studied and developed. In molecular classification techniques, the DNA barcode is a supporting tool for morphological classification (Hollingsworth et al., 2011). It is a modern technique that uses short DNA sequences to standardize differentiation between species (Kress, 2017). They have become a new tool to serve effectively in the inspection, classification, evaluation of genetic relationships, quality management, and origin of biological products (Mishra et al., 2015; Kress, 2017). In higher plants, some chloroplast genome regions (such as *matK*, *rbcl*, *psbA-trnH*, *atpF-atpH*, etc.) and nuclear genome regions (such as ITS-rDNA, 18S, etc.) are widely used in studying phylogenetic relationships, species taxonomy and identification (Kress et al., 2005; Mishra et al., 2015; Kress, 2017). However, for each different target group, the taxonomy of these genome regions was different (Kress et al., 2005; Tripathi et al., 2013).

Until now, the information about the DNA barcode of *T. petiolare* from Vietnam has not been studied. In this study, we carried out the identification of *T. petiolare* specimens on the basis of sequence analysis of 4 DNA barcode loci, namely ITS, *matK*, *rbcl* and *rpoC*. At the same time, this study also generated the database of DNA barcodes for *T. petiolare* plants collected in Vietnam for GenBank (NCBI..., 1988-2021).

Materials and methods

Plant materials

For this study, a total of 4 *Tinomiscium petiolare* samples were collected in Pu Luong Nature Reserve (PL1 and PL2) and Ben En National Park (BE1 and BE2) in Thanh Hoa province, Vietnam (Table 1). All these specimens were identified by Dr. Hoang Van Chinh (Hong Duc University) on the basis of their morphological characteristics. The specimens were deposited in the herbarium of Hong Duc University (Vietnam); 0.5 g of fresh leaves per plant sample were dried instantly in silica gel. All specimens were stored at -20°C until processed. Also, there were sequences downloaded from GenBank (Table 2).

DNA extraction, amplification and sequencing

Total DNA was extracted from leaf samples using DNeasy Plant Mini Kit (Qiagen, Germany). Amplification of the ITS, *matK*, *rbcl* and *rpoC* genes was carried out on the GeneAmp PCR System 9700 with the corresponding primers (Table 3). The PCR amplification reaction was performed in 25 µl reaction mixture consisting of 12.5 µl Master mix 2x (CWBIO, China), 0.75 µl each forward and reverse primer (10 pM/µl), 10 µl deionized water, and 1 µl of template DNA. The temperature cycle was as follows: one cycle of DNA denaturation at 94°C for 5 minutes followed by 35 cycles of 94°C for 1 minute, 52°C for 1 minute, and 72°C for 1 minute, with the last extension for 10 minutes at 72°C. The PCR product was tested on 0.8% agarose gel, then purified using GeneJET PCR Purification Kit (Thermo Fisher Scientific Co., USA). The sequence of DNA fragments was determined on the ABI 3500 Genetic Analyzer system following Sanger's principle, with BigDye® Terminator v3.1 Cycle Sequencing kit (Applied Biosystems Inc., USA) by sequencing directly from PCR products.

Sequence alignment and analysis

DNA sequences were analyzed using BioEdit software (Hall et al., 2011). All positions that contain gaps and missing data were removed from the data set. The sequences of 4 specimens were compared with those published on GenBank (Table 2) using the BLAST tool (McGinnis, Madden, 2004). The sequences were registered on GenBank (see Table 1 for the accession numbers). Also, there were sequences downloaded from GenBank (Table 2). The genetic distance and phylogenetic tree were calculated and constructed using MEGA6 software (Tamura et al., 2013).

Table 1. Specimens for testing potential barcodes, and accession numbers in GenBank

Таблица 1. Образцы растений для исследования потенциальных штрихкодов и номера образцов нуклеотидных последовательностей в базе данных GenBank

Voucher number	Species name	Location	GenBank accession number			
			ITS	<i>matK</i>	<i>rbcl</i>	<i>rpoC</i>
PL1	<i>Tinomiscium petiolare</i>	Pu Luong, Thanh Hoa, Vietnam	MW147627	MW123076	MW123080	MW123084
PL2	<i>Tinomiscium petiolare</i>	Pu Luong, Thanh Hoa, Vietnam	MW147628	MW123077	MW123081	MW123085
BE1	<i>Tinomiscium petiolare</i>	Ben En, Thanh Hoa, Vietnam	MW147629	MW123078	MW123082	MW123086
BE2	<i>Tinomiscium petiolare</i>	Ben En, Thanh Hoa, Vietnam	MW147630	MW123079	MW123083	MW123087

Table 2. Sequences from GenBank as extensions for species identification**Таблица 2. Последовательности из базы данных GenBank, использованные для видовой идентификации**

Species name	GenBank accession number			
	ITS	<i>matK</i>	<i>rbcL</i>	<i>rpoC</i>
<i>Tinomisium petiolare</i>	HG004877, KY365658	HG005005, DQ478612	KF181577, EF173675	
<i>Orthogynium</i> sp.	KY365652			
<i>Burasaia madagascariensis</i>	KY365641			
<i>Paratinospora sagittata</i>	KY365666, KY365671, KY365672, KY365673			
<i>Borismene japurensis</i>		KC494024		
<i>Burasaia apetala</i>		KC494025		
<i>Fibraurea tinctoria</i>		KC494035	HQ260781, FJ026485	
<i>Penianthus longifolius</i>		KC494046	FJ026499	
<i>Calycocarpum lyonii</i>		KC494026		
<i>Anamirta cocculus</i>		KC494022	FJ626591, EU526983	
<i>Sphenocentrum jollyanum</i>			JN051687	
<i>Tinospora sinensis</i>				MN727386
<i>Tinospora cordifolia</i>				NC042153, MH577056
<i>Sinomenium acutum</i>				MN626719
<i>Menispermum canadense</i>				NC048451, MH298221
<i>Menispermum dauricum</i>				NC042371, MH298220

Table 3. Primers used for amplification reactions in the study**Таблица 3. Праймеры, использованные для амплификации**

Locus	Primer name	Primer sequences (5'–3')	Expected product length (bp)	Reference
ITS	ITS1F	TCCGTAGGTGAACCTGCGG	650	White et al., 1990
	ITS4R	TCCTCCGTCTATTGATATGC		
<i>matK</i>	1RKIM-f	ACCCAGTCCATCTGGAAATCTTGGTTC	900	Ki-Joong Kim, pers. comm.
	3FKIM-r	CGTACAGTACTTTTGTGTTTACGAG		
<i>rbcL</i>	<i>rbcLa</i> -f	ATGTCACCACAAACAGAGACTAAAGC	600	Levin et al., 2003
	<i>rbcLa</i> -r	GTAATAATCAAGTCCACCRCG		Kress, Erickson, 2007
<i>rpoC</i>	<i>rpoCF</i>	TGAGAAAACATAAGTAAACGGGC	570	Ford et al., 2009
	<i>rpoCR</i>	GTGGATACACTTCTTGATATTGG		

Results and discussion

Total DNA extraction and amplification of gene segments

Total DNA products after extraction and purification were examined by electrophoresis on 0.8% agarose gel. It can be seen that the obtained DNA samples were of good quality. The electrophoresis clearly exhibited only a DNA tape with high molecular weight, sharpness, and quality assurance for the next study steps.

Total DNA of the specimens, after its ultraviolet absorbance (OD) had been determined by a spectrophotometer, was diluted to the concentration required to template PCR with the specific primers (Table 3). The results of the electrophoresis of PCR products on 0.8% agarose gel showed that the primers ITS, *matK*, *rbcl* and *rpoC* were used successfully for amplifying desired gene segments from DNA of 4 specimens and subsequent fragment cloning. Specifically, PCR products in 4 sequence regions ITS, *matK*, *rbcl* and *rpoC* of 4 studied specimens had sizes of about 650 bp, 950 bp, 600 bp and 600 bp, respectively (Fig. 1). This result is consistent with the sequences of primers, sufficiently reliable as a basis for reading ITS, *matK*, *rbcl* and *rpoC* sequences in the specimens to serve for the next studies.

Sequence determination

After purification of PCR products, we performed ITS, *matK*, *rbcl* and *rpoC* gene sequencing in 4 studied specimens on the ABI 3500 Genetic Analyzer, using BigDye® Terminator v3.1 Cycle Sequencing Kit. The results showed that all four specimens were completely similar in all 4 sequences of ITS, *matK*, *rbcl* and *rpoC*. The consensus ITS, *matK*, *rbcl* and *rpoC* sequences for the 4 studied specimens were deduced as follows:

ITS

GTCTGAATCGCAACCTTCTGGACGAGAGCCGGGCGGCCTCCGCCTTCCCCGGTGCTCGGCCGAAACAACAAACCCCGGCGCGGCACGCGCAAGGAAAACTCGAACGGAATTGGTGTGCCCGGACGATAGTGCATCGTCCCGGTGCCGCCGTCTCCGGGAAAAATCTCGAATGACTCTCGGCAACGGATATCTCGGCTCTCGCATCGATGAAGAACGTAGCGAAATGCGATACTTGGTGTGAATTGCAGAATCCCGTGAACCATCGAGTCTTTGAACGCAAGTTGCGCCC

GAACGCCGCTCCCGCCCCCTCGCAAGAGCGCGGACAGGAGCGAACGTTGGCCCCCGTGACCCGGCTCACGGTCGGCTTAAACGGATCCCCCTCGTTGCCCGCGACGCGATCGGTGGTGGTTGACGGCAACCCTTACCCGCGATTGGACGACGCGACCGAGGGGCACGGGGGAAACGAACCTTTCGGAGAGAACTCCACGAGCGACCTCAGGTCAGGCGGGGCCACCCGCTGAGTTTAA

matK

CTTTGCATTTATTGCGATTCTTTTCTACGAGTATCATAATTGGAATAGGCTTATTACTCAAAAAATAAATCCATTTCTGTTTTTCAAAAAAGAAAATCAAAGATTATTTCTGTTCCTATATAATTCTCATGTATATGAATGCGAATCATATTAGTTTTTTCCGTAAACAATCTTTTTATTACGATTAACTCTTCTAGAGCCTTTCTGGAGCGAACCCATTTCTATGAAAAATGGAACATCTTGTAGTAGTTTTTCAAAACGATTTTCAGTTTATCCTATGGTTGTTGAGGGAGCCTTTTCATGCATTATGTCAGATATCGAGGAAAAATCCATTCTGGGTTCAGGGGACCTTCTTCTGATGAATAAATGGAACATATTACCTGTAAATTTCTGGCAATGTAATTTTGACTTGTGGTCTCAACTGGATAGGATTTATATAACCCAATTAGCCAATCATTACTTTCGATTTTTTGGACTATCTTTCAAGTGTACGACTAAATACTTCGGTAGTAAGGAGTCAAATGTAGATAATTCAATTATTATGGATATTGCGATTAAGAAGTTCGATAGTATAGTTCCAATTATTTCTTGATTGGATCATTTGGCTAAAGCGAAATTTTGTAACGTATCAGGGCATCCCATTAGTAAGCCGGCTCGGGCCGATTTCATCAGATTCTGATATTATCGATAGATTTGGGCGAATATACAAAAATCTTTCTCATTATTACAGCGGATCCTCAAAAAAATAGTTTTGTATCGAATAAAGTATATACTTCGACTTTCCTGTGTGTAAGTCTTG

rbcl

AAGATTACAAATTGACTTATTATACTCCTGACTATGTACCCAAAGATACTGATACGCTAGCAGCATTCGGAGTAACTCCTCAACCTGGAGTTCCGCTGAAGAAGCGGGGCTGCGGTAGCTGCCGAATCTTCTACAGGTACATGGACAACGTGTGGACCGATGACTTACCAGTCTTGATCGTTACAAAGGACGATGCTACGACATTGAGCCCGTTGCTGGGGAAGAAAATCAATATATTGTTATGTAGCTTACCCCTTAGACCTTTTGAAGAAGGTTCTGTTACTATATGTTTACTTCCATTGTGGGTAATGTTTTTGGGTTCAAAGCGCTACGCGCTCTACGCTGGAGGATCTGCGAGTTCTACTGCTTATATTAACCTTTCCAAGGCCCGCTCATGGCATCCAAGTTGAGAGAGATAAATTGAACAAGTATGGTCTGCCCTATTGGGATGTACTATTAACCAAAATTTGGGATTATCCGCTAAGAAGTACGGTAGAGCAGTTTATGAATGTCTC




Fig. 1. DNA fragment amplification results for four studied specimens with primer pairs specific to the ITS, *matK*, *rbcl* and *rpoC* loci

Рис. 1. Результаты амплификации фрагментов ДНК четырех изученных образцов с парами праймеров, специфичных для локусов ITS, *matK*, *rbcl* и *rpoC*

rpoC

TTACAAGTCGTTTTTCAGATGTAATTGAAGGCAAA
GAGGGAAGATTTTCGCGAGACTCTGCTTGGCAAACGGGTC
GATTATTCGGGGCGTTCCGTCATTGTTGTGGGCCCTTC
GCTTTCATTAATCGATGTGGATTGCCTCGCGAAATAG
CAATAGAGCTTTTCAGACATTTGTCATTCTGTTGTTAAT
CAGACAACATATTGCTTCCAATATAGGGGTTGCTAAAAATA
AAATTCGGGAAAAAGAACCAATTGTGTGGGAAATAC
TTCAAGAAGTTATGCAGGGACATCCCGTATTGCTGAA
TAGAGCACCCACTCTGCATAGATTAGGCATACAGGCATTC
CAACCCATTTTAGTGAAGGACGTGCTATTTGTTTACATC
CATTAGTTTGTAAGGGATTCAATGCAGACTTTGATGGGGAT
CAAATGGCTGTTTCATGTACCTTTATCTTTGGAGGCT
CAAGCAGAGGCCCGTTTACTTATGTTTT

With 4 isolated sequences of ITS, *matK*, *rbcl* and *rpoC*, we identified 4 sequences of ITS, *matK*, *rbcl* and *rpoC* with the sizes 574 bp, 810 bp and 488 bp, respectively. Then, we analyzed and compared them with the ITS, *matK*, *rbcl* and *rpoC* sequences published in GenBank (NCBI <http://www.ncbi.nlm.nih.gov/>).

Sequence analysis

After obtaining the sequences of ITS, *matK*, *rbcl* and *rpoC*, we checked the similarity of the obtained sequences with those available in GenBank using the BLAST tool (McGinnis, Madden, 2004). In our study, the ITS, *matK* and *rbcl* sequences of the studied specimens showed a high similarity index corresponding to the ITS, *matK* and *rbcl* sequences of *T. petiolare* from GenBank. Specifically, the ITS sequence in the studied specimens showed a similarity index of 99.46–99.48% with the ITS sequence of *T. petiolare* from GenBank. The *matK* and *rbcl* sequences in the studied specimens showed a simi-

larity index of 100% with the *matK* and *rbcl* sequences of *T. petiolare* in GenBank. Meanwhile, when compared with the available sequences from GenBank, the *rpoC* sequence of the studied specimens showed a similarity index of 98.16% with *T. sinensis*. This discrepancy might be due to the fact that the *rpoC* sequence data for *T. petiolare* is not currently available in GenBank. The comparison with the GenBank database aimed to give a reference result with the group of species identical to the query nucleotide sequence. BLAST results could not lead to exact conclusions about species. In the cases of BLAST with high coverage and high homology (99%), it was not possible to revise the species name because BLAST results only showed the homogeneous nucleotide sequence that was available in GenBank. Since the results of BLAST showed inaccurate points, we conducted genetic distance determination and phylogenetic tree construction using MEGA6 software to determine the scientific names for the tested specimens (Tamura et al., 2013). The results are presented in Table 4 and Fig. 2.

The results in Table 4 demonstrate that the ITS, *matK*, *rbcl* and *rpoC* sequences of all the studied specimens collected in Ben En and Pu Luong (Thanh Hoa, Vietnam) had no differences. The ITS sequence of 4 studied specimens was not significantly different (0.13% and 0.4%) compared with the ITS sequence of *T. petiolare* published on GenBank with codes HG004877 and KY365658. Meanwhile, the difference between the ITS sequence that we obtained and the ITS sequences of some other species in the family Menispermaceae, published in GenBank with codes KY365652, KY365641, KY365673, KY365672, KY365671 and KY365666, was quite sizable. The *matK* and *rbcl* sequences produced from the studied specimens showed no difference when compared with *matK* and *rbcl* sequences of *T. petiolare* published

Table 4. Comparison of the studied gene sequences with the gene sequences in GenBank
(see Table 2 for the species name)

Таблица 4. Сравнительный анализ последовательностей изучаемых фрагментов и последовательностей, представленных в базе данных GenBank (см. видовые названия в таблице 2)


(a) ITS												
	BE1	BE2	PL1	PL2	HG004877	KY365658	KY365652	KY365641	KY365673	KY365672	KY365671	KY365666
BE1												
BE2	0.0000											
PL1	0.0000	0.0000										
PL2	0.0000	0.0000	0.0000									
HG004877	0.0013	0.0013	0.0013	0.0013								
KY365658	0.0040	0.0040	0.0040	0.0040	0.0013							
KY365652	0.1598	0.1598	0.1598	0.1598	0.1623	0.1715						
KY365641	0.1564	0.1564	0.1564	0.1564	0.1589	0.1679	0.0052					
KY365673	0.1645	0.1645	0.1645	0.1645	0.1671	0.1765	0.0200	0.0256				
KY365672	0.1645	0.1645	0.1645	0.1645	0.1671	0.1765	0.0200	0.0256	0.0000			
KY365671	0.1694	0.1694	0.1694	0.1694	0.1720	0.1818	0.0270	0.0327	0.0092	0.0092		
KY365666	0.1697	0.1697	0.1697	0.1697	0.1723	0.1820	0.0255	0.0298	0.0092	0.0092	0.0188	

Table 4. Continued
Таблица 4. Продолжение


(b) <i>matK</i>												
	BE1	BE2	PL1	PL2	HG005005	DQ478612	KC494024	KC494035	KC494025	KC494046	KC494026	KC494022
BE1												
BE2	0.0000											
PL1	0.0000	0.0000										
PL2	0.0000	0.0000	0.0000									
HG005005	0.0000	0.0000	0.0000	0.0000								
DQ478612	0.0000	0.0000	0.0000	0.0000	0.0000							
KC494024	0.0150	0.0150	0.0150	0.0150	0.0150	0.0156						
KC494035	0.0188	0.0188	0.0188	0.0188	0.0188	0.0195	0.0137					
KC494025	0.0188	0.0188	0.0188	0.0188	0.0188	0.0182	0.0125	0.0163				
KC494046	0.0126	0.0126	0.0126	0.0126	0.0126	0.0131	0.0075	0.0075	0.0088			
KC494026	0.0241	0.0241	0.0241	0.0241	0.0242	0.0250	0.0267	0.0280	0.0293	0.0192		
KC494022	0.0277	0.0277	0.0277	0.0277	0.0277	0.0274	0.0355	0.0381	0.0342	0.0293	0.0280	
(c) <i>rbcL</i>												
	BE1	BE2	PL1	PL2	KF181577	EF173675	FJ626591	JN051687	HQ260781	FJ026499	FJ026485	EU526983
BE1												
BE2	0.0000											
PL1	0.0000	0.0000										
PL2	0.0000	0.0000	0.0000									
KF181577	0.0000	0.0000	0.0000	0.0000								
EF173675	0.0000	0.0000	0.0000	0.0000	0.0000							
FJ626591	0.0057	0.0057	0.0057	0.0057	0.0057	0.0057						
JN051687	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0077					
HQ260781	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0076	0.0038				
FJ026499	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0077	0.0000	0.0038			
FJ026485	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0076	0.0038	0.0000	0.0038		
EU526983	0.0057	0.0057	0.0057	0.0057	0.0057	0.0057	0.0000	0.0077	0.0076	0.0077	0.0076	

Table 4. The end
Таблица 4. Окончание


(d) <i>rpoC</i>												
	BE1	BE2	PL1	PL2	MN727386	NC042153	MH577056	MN626719	NC048451	MH298221	NC042371	MH298220
BE1												
BE2	0.0000											
PL1	0.0000	0.0000										
PL2	0.0000	0.0000	0.0000									
MN727386	3.9708	3.9708	3.9708	3.9708								
NC042153	3.9704	3.9704	3.9704	3.9704	0.0021							
MH577056	3.9704	3.9704	3.9704	3.9704	0.0021	0.0000						
MN626719	3.9660	3.9660	3.9660	3.9660	0.0237	0.0259	0.0259					
NC048451	3.9656	3.9656	3.9656	3.9656	0.0259	0.0281	0.0281	0.0021				
MH298221	3.9656	3.9656	3.9656	3.9656	0.0259	0.0281	0.0281	0.0021	0.0000			
NC042371	3.9656	3.9656	3.9656	3.9656	0.0259	0.0281	0.0281	0.0021	0.0000	0.0000		
MH298220	3.9656	3.9656	3.9656	3.9656	0.0259	0.0281	0.0281	0.0021	0.0000	0.0000	0.0000	




(a) ITS tree



(b) *matK* tree



(c) *rbcL* tree



(d) *rpoC* tree

Fig. 2. Tree diagram showing the genetic relationship between the studied gene sequences and the gene sequences published in GenBank (see Table 2 for the species name)

Рис. 2. Дендрограмма, показывающая родство последовательностей изучаемых фрагментов и последовательностей, представленных в базе данных GenBank (см. видовые названия в таблице 2)

in GenBank with codes HG005005, DQ478612, KF181577 and EF173675. Meanwhile, the *matK* sequence that we obtained was different from the *matK* sequences of some other species in the same family Menispermaceae that were published in GenBank with codes KC494024, KC494035, KC494025, KC494046, KC494026 and KC494022. The difference rates were 1.50%, 1.88%, 1.88%, 1.26%, 2.41% and 2.77%, respectively. For the obtained *rbcL* sequence, there were differences with the *rbcL* sequences of some other species in the same family Menispermaceae published in GenBank with codes FJ626591, JN051687, HQ620781, FJ026499, FJ026485 and EU526983. The difference rates were 0.57%, 0.19%, 0.19%, 0.19%, and 0.57%, respectively. Notably, we did not find the *rpoC* sequence of *T. petiolare* on GenBank. So, the obtained *rpoC* sequence was significantly different from the *rpoC* sequences of other species in the same family Menispermaceae published in Genbank with codes MN727386, NC042153, MH577056, MN626719, NC048451, MH298221, NC042371 and MH298220. With these results, we confirmed that 4 studied specimens had genetic and species-specific relationships with *T. petiolare*.

Based on the identified sequences and homogenous ones from GenBank, we also built phylogenetic trees (Fig. 2). The results in Fig. 2 show that the ITS, *matK*, *rbcL* and *rpoC* sequences from the specimens that we collected and studied and the genetic sequence of *T. petiolare* were related to the same species and had a relatively high dissociation coefficient with other species. Therefore, the phylogenetic tree diagram demonstrates a diversity of branching, showing the genetic distance between the genetic sequence we obtained and the genetic sequences of some species that have been published in GenBank. The genetic sequences of the studied plant specimens collected at Pu Luong (PL1 and PL2) were not genetically different from those collected at Ben En (BE1 and BE2).

The results of the study have shown that the use of DNA barcodes is an effective method to quickly and accurately identify medicinal plant species (Chen et al., 2010; Mishra et al., 2016). Worldwide, there are also many studies that used DNA barcodes to identify species in the family Menispermaceae (Balasubramani, Venkatasubramanian, 2011; Yang et al., 2014; Osathanunkul et al., 2018; Wang et al., 2020). However, this study is the first to use a DNA barcode to authenticate *T. petiolare* from Vietnam. Of the four markers tested, ITS was arguably the most promising test region for species identification in this study. One advantage of the ITS region is that it can be amplified into two smaller segments (ITS1 and ITS2), therefore it is especially useful for degraded samples (Hillis, Dixon, 1991; Chen et al., 2010; Tripathi et al., 2013). The ITS region was also used to identify plant species with excellent results in previous studies (Balasubramani, Venkatasubramanian, 2011; Selvaraj et al., 2012; Thinh et al., 2020). At the same time, the results of this study also contribute to the DNA barcode database of *T. petiolare* in Vietnam.

Conclusions

We have succeeded in isolating and sequencing ITS, *matK*, *rbcL* and *rpoC* of the studied specimens collected at Pu Luong and Ben En (Thanh Hoa, Vietnam). The sequence sizes of ITS, *matK*, *rbcL* and *rpoC* that we obtained were 574 bp, 810 bp, 527 bp and 488 bp, respectively. The ITS, *matK*, *rbcL* and *rpoC* sequences of the studied specimens collected at Pu Luong and Ben En did not differ, showing that our specimens belonged to the same species group. The ITS, *matK*, *rbcL* and *rpoC* sequences of the studied specimens were identified as *T. petio-*

lare. For the first time, the *rpoC* sequence of *T. petiolare* was published in GenBank. The analysis also showed that there was still a lack of data on the genetic sequence of *T. petiolare* in other available databases than GenBank. Therefore, updating and supplementing molecular data about this species in order to have a sufficient basis for comparison and identification of plant samples is also a matter of concern and implementation.

The authors would like to thank Hong Duc University (Vietnam) and Far Eastern Federal University (Russia) for their support and facilitation to complete this study. The authors are also grateful to the Ministry of Science and Higher Education of the Russian Federation to provide Bui Bao Thinh a scholarship.

Авторы выражают благодарность Университету Хонгдык (Вьетнам) и Дальневосточному федеральному университету (Россия) за поддержку и помощь в осуществлении настоящего исследования. Кроме того, авторы благодарны Министерству науки и высшего образования Российской Федерации за предоставление стипендии Буи Бао Тхину.

References / Литература

- Balasubramani S.P., Venkatasubramanian P. Molecular identification and development of nuclear DNA ITS sequence-based marker to distinguish *Coscinium fenestratum* Gaertn. (Menispermaceae) from its adulterants. *Current Trends in Biotechnology and Pharmacy*. 2011;5(2):1163-1172.
- Chen S., Yao H., Han J., Liu C., Song J., Shi L. et al. Validation of the ITS2 region as a novel DNA barcode for identifying medicinal plant species. *PLoS One*. 2010;5(1):e8613. DOI: 10.1371/journal.pone.0008613
- Chi V.V. Dictionary of medicinal plants in Vietnam. Ho Chi Minh: Vietnam Publisher of Medicine; 2012. [in Vietnamese]
- Chinh V.T., Quang B.H., Anh T.T.P. Morphological characteristics and key to genera of family Menispermaceae in Vietnam. In: *Proceedings of the 6th National Scientific Conference of Ecology and Biological Resources (Hanoi, Vietnam)*; Hanoi; 2015. p.27-32.
- Christenhusz M.J., Byng J.W. The number of known plants species in the world and its annual increase. *Phytotaxa*. 2016;261(3):201-217. DOI: 10.11646/phytotaxa.261.3.1
- Ford C.S., Ayres K.L., Toomey N., Haider N., Van Alphen Stahl J., Kelly L.J. et al. Selection of candidate coding DNA barcoding regions for use on land plants. *Botanical Journal of the Linnean Society*. 2009;159(1):1-11. DOI: 10.1111/j.1095-8339.2008.00938.x
- Forman L.L. A synopsis of Thai Menispermaceae. *Kew Bulletin*. 1988;43(3):369-407. DOI: 10.2307/4118970
- Ghollasimood S., Faridah-Hanum I., Nazre M. Abundance and distribution of climbers in a coastal hill forest in Perak, Malaysia. *Journal of Agricultural Science*. 2012;4(5):245-254.
- Hall T., Biosciences I., Carlsbad C. BioEdit: an important software for molecular biology. *GERF Bulletin of Biosciences*. 2011;2(1):60-61.
- Hillis D.M., Dixon M.T. Ribosomal DNA: molecular evolution and phylogenetic inference. *The Quarterly Review of Biology*. 1991;66(4):411-453. DOI: 10.1086/417338

- Ho P.H. An illustrated flora of Vietnam. Vol. 1. Ho Chi Minh, Vietnam: Young Publishing House; 2000. [in Vietnamese]
- Hollingsworth P.M., Graham S.W., Little D.P. Choosing and using a plant DNA barcode. *PLoS One*. 2011;6(5):e19254. DOI: 10.1371/journal.pone.0019254
- Kress W.J. Plant DNA barcodes: Applications today and in the future. *Journal of Systematics and Evolution*. 2017;55(4):291-307. DOI: 10.1111/jse.12254
- Kress W.J., Erickson D.L. A two-locus global DNA barcode for land plants: the coding *rbcl* gene complements the non-coding *trnH-psbA* spacer region. *PLoS One*. 2007;2(6):e508. DOI: 10.1371/journal.pone.0000508
- Kress W.J., Wurdack K.J., Zimmer E.A., Weigt L.A., Janzen D.H. Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;102(23):8369-8374. DOI: 10.1073/pnas.0503123102
- Levin R.A., Wagner W.L., Hoch P.C., Nepokroeff M., Pires J.C., Zimmer E.A. et al. Family-level relationships of Onagraceae based on chloroplast *rbcl* and *ndhF* data. *American Journal of Botany*. 2003;90(1):107-115. DOI: 10.3732/ajb.90.1.107
- McGinnis S., Madden T.L. BLAST: at the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Research*. 2004;32 Suppl 2:W20-W25. DOI: 10.1093/nar/gkh435
- Mishra P., Kumar A., Nagireddy A., Mani D.N., Shukla A.K., Tiwari R. et al. DNA barcoding: an efficient tool to overcome authentication challenges in the herbal market. *Plant Biotechnology Journal*. 2016;14(1):8-21. DOI: 10.1111/pbi.12419
- NCBI. National Center for Biotechnology Information. U.S. National Library of Medicine. Bethesda, MD; 1988-2021. Available from: <http://www.ncbi.nlm.nih.gov/> [accessed Aug. 31, 2020].
- Osathanunkul M., Osathanunkul R., Madesis P. Species identification approach for both raw materials and end products of herbal supplements from *Tinospora* species. *BMC Complementary and Alternative Medicine*. 2018;18(1):111. DOI: 10.1186/s12906-018-2174-0
- Selvaraj D., Shanmughanandhan D., Sarma R.K., Joseph J.C., Srinivasan R.V., Ramalingam S. DNA barcode ITS effectively distinguishes the medicinal plant *Boerhavia diffusa* from its adulterants. *Genomics, Proteomics and Bioinformatics*. 2012;10(6):364-367. DOI: 10.1016/j.gpb.2012.03.002
- Tamura K., Stecher G., Peterson D., Filipski A., Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*. 2013;30(12):2725-2729. DOI: 10.1093/molbev/mst197
- Thinh B.B., Chac L.D., Thu L.T. Application of internal transcribed spacer (ITS) sequences for identifying *Anoectochilus setaceus* Blume in Thanh Hoa, Vietnam. *Proceedings on Applied Botany, Genetics and Breeding*. 2020;181(2):108-116. DOI: 10.30901/2227-8834-2020-2-108-116
- Tripathi A.M., Tyagi A., Kumar A., Singh A., Singh S., Chaudhary L. et al. The internal transcribed spacer (ITS) region and *trnH-psbA* are suitable candidate loci for DNA barcoding of tropical tree species of India. *PLoS One*. 2013;8(2):e57934. DOI: 10.1371/journal.pone.0057934
- Van Valkenburg J.L.C.H., Bunyapraphatsara N. Medicinal and poisonous plants 2. In: *Plant Resources of South-East Asia*. Vol 12(2). Leiden: Backhuys Publishers; 2001. p.550-552.
- Wang X., Xue J., Zhang Y., Xie H., Wang Y., Weng W. et al. DNA barcodes for the identification of *Stephania* (Menispermaceae) species. *Molecular Biology Reports*. 2020;47(3):2197-2203. DOI: 10.1007/s11033-020-05325-6
- White T.J., Bruns T., Lee S., Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: M.A. Innis, D.H. Gelfand, J.J. Sninsky, T.J. White (eds). *PCR Protocols: A Guide to Methods and Applications*. London: Academic Press; 1990. p.315-322.
- Yang P., Li X., Zhou H., Hu H., Zhang H., Sun W., et al. Molecular identification of Chinese materia medica and its adulterants using ITS2 and *psbA-trnH* Barcodes: a case study on Rhizoma Menisperm. *Chinese Medicine*. 2014;5(4):1-8. DOI: 10.4236/cm.2014.54023

Прозрачность финансовой деятельности / The transparency of financial activities

Авторы не имеют финансовой заинтересованности в представленных материалах или методах.

The authors declare the absence of any financial interest in the materials or methods presented.

Для цитирования / How to cite this article

Тхинь Б.Б., Дудкин Р.В., Чак Л.Д., Чинь Х.В., Хой К.В., Лиен Н.Т. Идентификация *Tinomisium petiolare* из Вьетнама с помощью штрих-кода ДНК. Труды по прикладной ботанике, генетике и селекции. 2021;182(2):114-122. DOI: 10.30901/2227-8834-2021-2-114-122

Thinh B.B., Doudkin R.V., Chac L.D., Chinh H.V., Hoi Q.V., Lien N.T. Identification of *Tinomisium petiolare* from Vietnam using the DNA barcode. *Proceedings on Applied Botany, Genetics and Breeding*. 2021;182(2):114-122. DOI: 10.30901/2227-8834-2021-2-114-122

Авторы благодарят рецензентов за их вклад в экспертную оценку этой работы / The authors thank the reviewers for their contribution to the peer review of this work

Дополнительная информация / Additional information

Полные данные этой статьи доступны / Extended data is available for this paper at <https://doi.org/10.30901/2227-8834-2021-2-114-122>

Мнение журнала нейтрально к изложенным материалам, авторам и их месту работы / The journal's opinion is neutral to the presented materials, the authors, and their employer

Авторы одобрили рукопись / The authors approved the manuscript

Конфликт интересов отсутствует / No conflict of interest

ORCID

Thinh B.B. <https://orcid.org/0000-0002-3826-1199>
Doudkin R.V. <https://orcid.org/0000-0002-5528-7729>
Chac L.D. <https://orcid.org/0000-0002-0845-4678>

Chinh H.V. <https://orcid.org/0000-0002-5329-9478>
Hoi Q.V. <https://orcid.org/0000-0002-3691-941X>
Lien N.T. <https://orcid.org/0000-0002-0069-9945>